



Crop Management

Considerations for Selecting Soybean Varieties

Selecting soybean varieties is one of the most critical decisions a farmer can make each growing season. Although high yield potential is important, there are also other factors you should consider when selecting varieties, including disease, insect, and herbicide resistance traits; maturity date; and target market.

This publication describes the main factors you should consider when selecting which soybean variety to plant. Here is a quick list of considerations for selecting soybean varieties:

- Choose varieties that perform well in multi-location performance trials. Do not rely solely on data from your farm or one location or source to make decisions.
- Select a range of maturity groups and varieties that best match your production practices.
- Pick varieties that yield well and have the traits that are important to you (such as herbicide or disease resistance).
- Consider field history when selecting varieties, particularly when you look for varieties with disease and insect resistance.
- Select different varieties to diversify genetics. Plant the majority of your acreage to proven varieties and test new varieties on a smaller scale.
- Pay attention to seed composition characteristics (for example, protein, oil, amino acids, and so on). These qualities may not affect your “premium” at the local elevator, but they can affect exports.
- Select varieties with low lodging potential for planting into irrigated fields or mucky fields. Lodging can increase harvest losses and significantly delay harvest operations.
- Select varieties with high ratings for tolerance to iron deficiency chlorosis (IDC). IDC can reduce yield potential in some areas.
- Buy only the traits you need. Pest management traits protect yield, they do not enhance it.



Figure 1. Iron deficiency chlorosis can cause yield loss in certain regions of the United States.

Herbicide Traits

It is important to know the weed history, previous herbicide use patterns, and herbicide-resistant weed issues in a particular field to help you choose the herbicide resistance traits you need when choosing a soybean variety.

There are many varieties available with and without herbicide resistance traits. The most common herbicide resistance traits include:

- Conventional (non-traited)
- Roundup Ready®, which allows for in-season glyphosate application
- LibertyLink®, which allows for in-season glufosinate application
- Roundup Ready 2 Xtend®, which allows for at-planting and in-season applications of dicamba and glyphosate
- Enlist®, which allows for at-planting and in-season applications of 2, 4-D choline, glyphosate, and glufosinate — note that international market approval for this trait is pending
- BOLT® soybean, which has enhanced tolerance to certain ALS inhibitor-type herbicides, and in-season glyphosate applications

Disease Considerations

Planting disease-resistant soybean varieties can be the most effective and economical disease management option. Soybean varieties are available that have varying levels of resistance to some of the major diseases that occur in the United States and Ontario, Canada, such as soybean cyst nematode, sudden death syndrome, and frogeye leaf spot.

Some varieties may include resistance ratings for other diseases, but not all varieties are rated for every disease. If you have a field with a history of one or more diseases, select varieties that are resistant to the diseases that have the greatest effect on yield.

Insect Considerations

There are resistance traits available that protect soybean against soybean aphid in several maturity groups. Soybean aphid-resistant varieties have *Rag1*, *Rag2*, or both resistance genes. The *Rag* resistance genes are non-GMO, so they are available for organic farmers.

The maturity group you select can also help mitigate the damage from some insect pests. For example, early-maturing varieties are often less susceptible to late-season pod feeding by bean leaf beetles or stink bugs.

Maturity

Several factors affect soybean development, including genetics, temperature, and hours of sunshine. Plus, diseases, moisture, and other stresses can lengthen or shorten the actual days to maturity depending on when the stress occurs. For these reasons, it's important to routinely evaluate the ability of new soybean varieties to adapt to these conditions. After this evaluation, choose the most appropriate maturity group (MG) to maximize yields and returns.




Figure 2. Optimum soybean maturity across the United States. Source: www.coolbean.info/library/documents/SoybeanMG_2016_FINAL.pdf.

The Bottom Line

Variety selection is key to maximizing yield potential. Carefully consider multiple factors when selecting varieties.

Other publications in the *Crop Management* series are available on the Crop Protection Network website (CropProtectionNetwork.org).

In particular, see: *Delineating Optimal Soybean Maturity Groups Across the United States* from University of Wisconsin Extension at www.coolbean.info/library/documents/SoybeanMG_2016_FINAL.pdf.

Authors

Shawn Conley, University of Wisconsin-Madison
 Carl Bradley, University of Kentucky
 Martin Chilvers, Michigan State University
 Loren Giesler, University of Nebraska
 Daren Mueller, Iowa State University
 Ed Sikora, Auburn University
 Damon Smith, University of Wisconsin-Madison
 Albert Tenuta, OMAFRA
 Kelley Tilmon, Ohio State University
 Kiersten Wise, University of Kentucky



Reviewers

Tom Allen, Mississippi State University
 Alyssa Collins, Pennsylvania State University
 Nick Dufault, University of Florida
 Travis Faske, University of Arkansas
 J.D. Green, University of Kentucky
 Louis Hesler, USDA-ARS-NCARL
 Erin Hodgson, Iowa State University
 Doug Jardine, Kansas State University
 Bill Johnson, Purdue University
 Heather Kelly, University of Tennessee
 Jan Knodel, North Dakota State University
 Christian Krupke, Purdue University
 Travis Legleiter, University of Kentucky
 Laura Linney, Ohio State University
 Dean Malvick, University of Minnesota
 Sam Markell, North Dakota State University
 Brian McCornack, Kansas State University
 Hillary Mehl, Virginia Tech
 Bruce Potter, University of Minnesota
 Deirdre Prischmann-Voldseth, North Dakota State University
 Adam Sisson, Iowa State University
 Scott Stewart, University of Tennessee
 Lindsey Thiessen, North Carolina State University

Photo Credits

All photos were provided by and are the property of the authors and reviewers, except Figure 1 provided by Greg Tylka, Iowa State University.

Acknowledgments

The *Crop Management* series is a multi-state collaboration sponsored by the North Central Soybean Research Program (NCSRP, www.ncsrp.com). This publication was developed by the Crop Protection

Network, a multi-state and international collaboration of university/provincial extension specialists and public/private professionals that provides unbiased, research-based information to farmers and agricultural personnel.

The authors thank the United Soybean Board, USDA-NIFA, and Grain Farmers of Ontario for their support.

Design and production by Purdue Agricultural Communication

The information in this publication is only a guide, and the authors assume no liability for practices implemented based on this information. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.



The information in this publication is only a guide, and the authors assume no liability for practices implemented based on this information. Reference to products in this publication is not intended to be an endorsement to the exclusion of others that may be similar. Individuals using such products assume responsibility for their use in accordance with current directions of the manufacturer.

Feb 2018

The U.S. Department of Agriculture (USDA) prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex, marital status, familial status, parental status, religion, sexual orientation, genetic information, political beliefs, reprisal, or because all or a part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) Persons with disabilities who require alternative means for communication of program information (Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD). To file a complaint of discrimination write to USDA, Director, Office of Civil Rights, 1400 Independence Avenue, S.W., Washington, D.C. 20250-9410 or call (800) 795-3272 (voice) or (202) 720-6382 (TDD). USDA is an equal opportunity provider and employer.

©2018 by the Crop Protection Network. All rights reserved.